Progress Report

Matt Gordon

August 10, 2018

Geometric Aberration Study

- Simple Test Case: Use emittance component tool to identify geometric aberration emittance growth in solenoid
- Using simple current loop, calculating emittance growth due to spherical aberration from:
 - Kumar, Vinit & Phadte, Deepraj & Bhai Patidar, Chirag. (2011). A simple formula for emittance growth due to spherical aberration in a solenoid lens.
- Compare to simulation

Spherical Aberration Calculation

 The emittance growth of a azimuthally symmetric beam due to a solenoid in the thin lens approximation is simply related to the geometry of the solenoid and the beam


$$\varepsilon_{x y} = \frac{R^4}{2\sqrt{6} f_0} \sqrt{\frac{C_1^2}{12} + \frac{C_1 C_2}{5} R^2 + \frac{C_2^2}{8} R^4}$$

Where C1 and C2 are reductions of the focal length due to the 3rd and 5th order spherical aberrations respectively

$$C_{1} = \frac{1}{2} \frac{\int_{-\infty}^{+\infty} \{B'(z)\}^{2} dz}{\int_{-\infty}^{+\infty} B^{2}(z) dz}, \quad C_{2} = \frac{5}{64} \frac{\int_{-\infty}^{+\infty} \{B''(z)\}^{2} dz}{\int_{-\infty}^{+\infty} B^{2}(z) dz}.$$


Comparison


 Started by getting unreasonable results as compared to paper, several orders of magnitude larger (10^-9 compared to 10^-11)



Time Step Convergence

• Tested in case where emittance growth should be negligible (10^-15)Trouble converging in reasonable time step run time (~1 hours)





Comparing Emittance Component to Emittance (~10 hour run time)

- Expected emittance growth from spherical aberration (2*10^-10 m)
- Doesn't seem to line

