


Initial CLEO-c Results

Anders Ryd
Cornell University



Nov. 29, 2005

$$e^+ e^- \rightarrow c \bar{c} \rightarrow D^0 \bar{D}^0$$
$$\bar{D}^0 \rightarrow K^+ \pi^-, D^0 \rightarrow K^- e^+ \bar{\nu}$$




Testing the Quark Mixing (CKM) Matrix

- The CKM matrix provides the only mechanism for CP violation in the SM.
- It is an important goal of flavor physics to measure – and overconstrain – the parameters in the CKM matrix.
- Non-perturbative strong effects limit our ability to extract the fundamental parameters from the measurements.
- **CLEO-c provides unique measurements that will address this limitation.**




Testing Theories of Strong Interactions




- Measure form factors in $D \rightarrow \pi l \nu$ and validate theoretical calculations
 - Can then use this to extract $|V_{ub}|$ from $B \rightarrow \pi l \nu$

- B mixing is well measured
 - $\Delta m_d = (0.502 \pm 0.007) \times 10^{-12} \text{ s}^{-1}$
- But $|V_{td}|$ from Δm_d has large uncertainties from f_B
- CLEO-c can measure f_D



$$\Delta m_d = \frac{G_F^2}{6} M_B M_t^2 |V_{td} V_{tb}^*|^2 \eta_B S_0(x_t) f_B^2 B_B$$



Outline

- CLEO-c Program
- Experimental technique
- Hadronic branching fractions
 - Reference branching fractions
 - Cabibbo suppressed decays
- Semileptonic decays
 - Inclusive branching fractions
 - Exclusive branching fractions
 - Form factors
- Leptonic decays
- D_s scan
- Conclusions

Normalization Br. Fr.
Systematics



Tests of theory for
strong physics:
 f_D , form factors

CLEO-c Program

- The CLEO-c program as proposed (CLNS 01/1742)
- Three year program
 - 1 year (3 fb^{-1}) at $\psi(3770)$ for $D\bar{D}$ physics
 - 1 year (3 fb^{-1}) above $D_s D_s$ threshold for D_s physics
 - 1 year at J/ψ for glueballs, exotics etc.
- Program was approved for running in 2003 until Mar 31, 2008.

CESR-c

- CESR was upgraded with wiggler magnets to increase damping at lower energies
- These super conducting magnets have worked very well
- However, the currents that can be stored are much lower than planned.
 - Luminosity about $\frac{1}{4}$ of design
- Will install compensating solenoids to gain about 50%




CLEO-c Plans

- With a better understanding of the performance of CESR-c and looking at the physics case we are now looking at a somewhat modified run plan
 - $\sim 750 \text{ pb}^{-1}$ at $\psi(3770)$
 - $\sim 750 \text{ pb}^{-1}$ at $D_s D_s$ threshold
 - $\sim 5\%$ of running time at $\psi(2S)$, 30M events
- Solid motivation for J/ψ running was hard to find. The $f_J(2220)$ no longer there. Studying this was an important goal for the J/ψ program.
- We have done a scan in the D_s threshold region to understand where to run and measure the cross-section

CLEO-c detector


2230104-002

CLEO-c



- 1 T B-field.
- New 6-layer inner drift chamber.
- Otherwise the CLEO III detector

ZD: New Inner Drift Chamber




- Replace CLEO III silicon vertex detector.
- Less material – important for the lower momentum particles at charm threshold.
- Detector was installed in summer '03 and is working well.

Physics at $c\bar{c}$ Threshold

- We run at $E_{\text{cm}} = 3.77$ GeV
 - the $\psi(3770)$ resonance.
- Producing $D\bar{D}$ pairs
 - and no other pions.
- Makes this a very clean experiment for studies of charm decays.
- Most analyses use a tagging technique
 - one of the produced D mesons is fully reconstructed.

$$e^+ e^- \rightarrow c \bar{c} \rightarrow D^0 \bar{D}^0$$

$$\bar{D}^0 \rightarrow K^+ \pi^-, D^0 \rightarrow K^- e^+ \bar{\nu}$$



Data Samples at $\psi(3770)$

- Analyses presented are based on two data samples
 - 56 pb⁻¹ recorded Dec '03 - Apr '04
 - 281 pb⁻¹ recorded Dec '03 – Apr '05 (include the 56 pb⁻¹)
 - We project that we will have 750 pb⁻¹ by April 2008.
- We have final results from 281 pb⁻¹ for $D^+ \rightarrow \mu^+ \nu_\mu$
- Many other analyses are being finalized
 - Systematics studies need to be done in detail for many analyses

Tag Reconstruction

- Fully reconstructed D -tag candidates are built from
 - Charged kaons, good tracks from IP with particle identification
 - Charged pions, good tracks from IP with particle identification
 - π^0 ($\gamma\gamma$)
 - K_s ($\pi^+\pi^-$), no flight significance cut applied.
- To extract yields we typically cut on $\Delta E = E_{\text{Reco}} - E_{\text{beam}}$
 - Cut is mode (and analysis) dependent
- and fit the m_{BC} distribution
- We reconstruct a D^0 tag with $\sim 17\%$ eff. and D^+ tag with $\sim 10\%$ eff.
 - Different analyses use different tag modes.

Hadronic D -decays

- Hadronic decays of D mesons are important as they set the branching fraction scale of many B and D decays.
- The PDG '04 is dominated by measurements based on the tagging of a D^* by the presence of a slow pion (CLEO and LEP experiments)
 - D^+ branching fractions are less well known, as a slow π^0 is not a clean tag.
 - Systematics limited.
 - Our systematics are orthogonal.
- As the double tag technique we use determines the number of $D\bar{D}$ pairs we can also calculate the cross-section for D^+D^- and $D^0\bar{D}^0$ production.

Hadronic D -decays and $\sigma(e^+e^- \rightarrow D\bar{D})$

- In order to measure the cross section and absolute branching fractions we need to determine the number of produced $D\bar{D}$ events
 - Use a 'double tag' technique, pioneered by MARK III

$$N_i = 2 \epsilon_i B_i N_{D\bar{D}}$$

$$N_{ij} = \epsilon_{ij} B_i B_j N_{D\bar{D}}$$

$$N_{D\bar{D}} = \frac{N_i N_j}{4 N_{ij}} \frac{\epsilon_{ij}}{\epsilon_i \epsilon_j}$$

- Use 3 D^0 modes ($K^-\pi^+$, $K^-\pi^+\pi^0$, and $K^-\pi^+\pi^-\pi^+$) and 6 D^+ modes ($K^-\pi^+\pi^+$, $K_s^-\pi^+$, $K^-\pi^+\pi^+\pi^0$, $K_s^-\pi^+\pi^-\pi^+$, $K_s^-\pi^+\pi^0$, and $K^-\pi^+\pi^+$)
- Determine separately the D and \bar{D} yields
 - This gives 18 single tag yields and 45 ($=3^2+6^2$) double tag yields
- In a combined χ^2 fit we extract 9 branching fractions and $D^0\bar{D}^0$ and D^+D^- yields. The fit includes the systematic errors.
- Many systematics cancel in the $D\bar{D}$ yield (e.g. tracking eff, PID eff.).

Initial State Radiation

- We run at $E_{cm} = 3.77$ GeV to produce the $\psi(3770)$
 - The spread in E_{cm} is about 2 MeV
 - The width of the $\psi(3770)$ is about 25 MeV
- However, the beam particles can radiate a photon and produce the $\psi(3770)$ at a lower energy.
 - In fact in every interaction many photons are emitted, but at such a low energy that we can not detect them. The distribution of energy radiated by (soft) photons is given by:

$$f(E_\gamma) \propto E_\gamma^{\beta-1}$$
$$\beta = \frac{2\alpha}{\pi} \left[2 \ln \frac{E_{cm}}{m_e} - 1 \right] \approx 0.07$$

- Many analyses use a first principle lineshape for the m_{BC} fit

ISR in Data vs. MC





D⁺ → K⁻ π⁺ π⁺

DATA

D⁺ → K⁻ π⁺ π⁺



Monte Carlo



$$M_{BC} = \sqrt{E_{\text{beam}}^2 - |p(D)|^2}$$


Single Tag Yields (56 pb⁻¹)

D or \bar{D} Mode	Yield (10^3)	Efficiency (%)
$D^0 \rightarrow K^- \pi^+$	5.11 ± 0.07	64.6 ± 0.3
$\bar{D}^0 \rightarrow K^+ \pi^-$	5.15 ± 0.07	65.6 ± 0.3
$D^0 \rightarrow K^- \pi^+ \pi^0$	9.51 ± 0.11	31.4 ± 0.1
$\bar{D}^0 \rightarrow K^+ \pi^- \pi^0$	9.47 ± 0.11	31.8 ± 0.1
$D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$	7.44 ± 0.09	43.6 ± 0.2
$\bar{D}^0 \rightarrow K^+ \pi^- \pi^- \pi^+$	7.43 ± 0.09	43.9 ± 0.2
$D^+ \rightarrow K^- \pi^+ \pi^+$	7.56 ± 0.09	50.7 ± 0.2
$D^- \rightarrow K^+ \pi^- \pi^-$	7.56 ± 0.09	51.3 ± 0.2
$D^+ \rightarrow K^- \pi^+ \pi^+ \pi^0$	2.45 ± 0.07	25.7 ± 0.2
$D^- \rightarrow K^+ \pi^- \pi^- \pi^0$	2.39 ± 0.07	25.7 ± 0.2
$D^+ \rightarrow K_S^0 \pi^+$	1.10 ± 0.04	45.5 ± 0.4
$D^- \rightarrow K_S^0 \pi^-$	1.13 ± 0.04	45.9 ± 0.4
$D^+ \rightarrow K_S^0 \pi^+ \pi^0$	2.59 ± 0.07	22.4 ± 0.2
$D^- \rightarrow K_S^0 \pi^- \pi^0$	2.50 ± 0.07	22.4 ± 0.2
$D^+ \rightarrow K_S^0 \pi^+ \pi^+ \pi^-$	1.63 ± 0.06	31.1 ± 0.2
$D^- \rightarrow K_S^0 \pi^- \pi^- \pi^+$	1.58 ± 0.06	31.3 ± 0.2
$D^+ \rightarrow K^+ K^- \pi^+$	0.64 ± 0.03	41.4 ± 0.5
$D^- \rightarrow K^+ K^- \pi^-$	0.61 ± 0.03	40.8 ± 0.5




Single Tag Yields (56 pb⁻¹)

3970305-001



Double Tag Fits




- Beam energy spread and ISR causes a correlated shift in the mass of the two D -mesons
- Resolution is uncorrelated among the two D -mesons
- A **two dimensional fit** allow us to separate the effects of beam energy smearing and detector resolution.

MC $\sigma_E = (2.08 \pm 0.01) \text{ MeV}$

Data $\sigma_E = (2.11 \pm 0.02) \text{ MeV}$

- MC width comes from machine parameters

Double Tag Yields (56 pb^{-1})



- The statistical errors on the double tag yields set the errors on the branching fractions (assuming the single tag yields don't dominate the errors).

Results 56 pb⁻¹

Parameter	Fitted Value	Δ_{FSR}	PDG 2004*
$N_{D^0\bar{D}^0}$	$(2.01 \pm 0.04 \pm 0.02) \times 10^5$	-0.2%	
$\mathcal{B}(D^0 \rightarrow K^- \pi^+)$	$(3.91 \pm 0.08 \pm 0.09)\%$	-2.0%	$\epsilon=(95.43 \pm 0.04)\%$
$\mathcal{B}(D^0 \rightarrow K^- \pi^+ \pi^0)$	$(14.9 \pm 0.3 \pm 0.5)\%$	-0.8%	
$\mathcal{B}(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)$	$(8.3 \pm 0.2 \pm 0.3)\%$	-1.7%	
$N_{D^+D^-}$	$(1.56 \pm 0.04 \pm 0.01) \times 10^5$	-0.2%	
$\mathcal{B}(D^+ \rightarrow K^- \pi^+ \pi^+)$	$(9.5 \pm 0.2 \pm 0.3)\%$	-2.2%	$(9.2 \pm 0.6)\%$
$\mathcal{B}(D^+ \rightarrow K^- \pi^+ \pi^+ \pi^0)$	$(6.0 \pm 0.2 \pm 0.2)\%$	-0.6%	
$\mathcal{B}(D^+ \rightarrow K_S^0 \pi^+)$	$(1.55 \pm 0.05 \pm 0.06)\%$	-1.8%	
$\mathcal{B}(D^+ \rightarrow K_S^0 \pi^+ \pi^0)$	$(7.2 \pm 0.2 \pm 0.4)\%$	-0.8%	
$\mathcal{B}(D^+ \rightarrow K_S^0 \pi^+ \pi^+ \pi^-)$	$(3.2 \pm 0.1 \pm 0.2)\%$	-1.4%	
$\mathcal{B}(D^+ \rightarrow K^+ K^- \pi^+)$	$(0.97 \pm 0.04 \pm 0.04)\%$	-0.9%	

*Our branching fractions are corrected for FSR, PDG values are not


Using our measured luminosity of $55.8 \pm 0.6 \text{ pb}^{-1}$ we obtain:

$$\sigma(D^0 \bar{D}^0) = (3.60 \pm 0.07 \pm 0.07) \text{ nb} \quad \sigma(D^+ D^-) = (2.79 \pm 0.07 \pm 0.10) \text{ nb}$$


$$\sigma(D \bar{D}) = (6.39 \pm 0.10 \pm 0.17) \text{ nb}$$

Comparison with Other Exp.


$$Br(D^0 \rightarrow K^- \pi^+)$$



$$Br(D^+ \rightarrow K^- \pi^+ \pi^+)$$





Comparison to PDG



Projections

- These results were based on 56 pb^{-1} (PRL 95, 121801 (2005))
- We are now analyzing the 281 pb^{-1} sample



- Statistical uncertainties about 1% in the 281 pb^{-1} sample!



$D^0 \rightarrow K^- \pi^+$ and $D^+ \rightarrow K^- \pi^+ \pi^+$ Systematics

Source	56 pb ⁻¹	281 pb ⁻¹	750 pb ⁻¹
Trk. eff.	0.7%	0.35%	0.25%
Kaon PID eff.	1.3%	0.3%	0.3%
Pion PID eff.	0.3%	0.2%	0.2%
ΔE selection	1.0%	0.3%	0.3%
Fit shape	0.8%	0.5%	0.5%
FSR	0.5%	0.5%	0.5%
Res. Sub. Structure	0.6%	0.4%	0.4%
Double DCSD interf.	0.8%	0.8%	0.5%
<hr/>			
$D^0 \rightarrow K^- \pi^+$ Stat	2.1%	0.9%	0.6%
$D^0 \rightarrow K^- \pi^+$ Syst	3.1%	1.3%	1.1%
$D^0 \rightarrow K^- \pi^+$ Total	3.7%	1.6%	1.2%
<hr/>			
$D^+ \rightarrow K^- \pi^+ \pi^+$ Stat	3.9%	1.2%	0.7%
$D^+ \rightarrow K^- \pi^+ \pi^+$ Syst	2.6%	1.4%	1.2%
$D^+ \rightarrow K^- \pi^+ \pi^+$ Total	4.7%	1.8%	1.4%




- Systematics limited at 281 pb⁻¹
 - Some systematics should improve with additional statistics

Tracking Efficiencies

- Events that can be fully reconstructed can be used for very clean studies of tracking efficiencies.
- We have used DD events and $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$ events
- Look at recoil mass against D^0 -tag and pion – see how often kaon is found
 - In data we find $\epsilon = (90.8 \pm 0.4)\%$



$D^+ \rightarrow K_{L,S} \pi^+$



Interfere destructively
(Long D^+ lifetime)

$$\bar{K}^0 = \frac{1}{\sqrt{2}} (K_S^0 - K_L^0)$$

$$K^0 = \frac{1}{\sqrt{2}} (K_S^0 + K_L^0)$$


The physical states of the K_S and K_L have different rates due to interference

Based on factorization Bigi and Yamamoto (PLB 349, 363 (1995))
 Predicts


$$\frac{\Gamma(D^+ \rightarrow K_L) - \Gamma(D^+ \rightarrow K_S)}{\Gamma(D^+ \rightarrow K_L) + \Gamma(D^+ \rightarrow K_S)} \approx 10\%$$

Preliminary Results

281 fb⁻¹



Reconstruct tag D and pion, look for signal in the recoil mass




- $\mathcal{B}(D^+ \rightarrow K^0_s \pi^+) + \mathcal{B}(D^+ \rightarrow K^0_L \pi^+) = (3.06 \pm 0.06 \pm 0.16)\%$
- $\text{Asymmetry} = (K^0_L - K^0_s)/(K^0_L + K^0_s) = -0.01 \pm 0.04 \pm 0.07$
 - Consistent with 10% prediction.
- $\mathcal{B}(D^+ \rightarrow \eta \pi^+) = (0.39 \pm 0.03 \pm 0.03)\%$ [PDG2004 has $(0.30 \pm 0.06)\%$].

$D \rightarrow n(\pi^\pm)m(\pi^0)$ (Preliminary)

- This analysis doesn't use D -tags.
- Measure ratio to normalization mode

Mode	$\mathcal{B} (\times 10^{-3})$	PDG ($\times 10^{-3}$)
$\pi^+\pi^-$	$1.40 \pm 0.04 \pm 0.03$	1.38 ± 0.05
$\pi^0\pi^0$	$0.78 \pm 0.05 \pm 0.04$	0.84 ± 0.22
$\pi^+\pi^-\pi^0$	$13.3 \pm 0.2 \pm 0.5$	11 ± 4
$\pi^0\pi^0\pi^0$	< 0.30	---
$\pi^+\pi^+\pi^-\pi^-$	$7.42 \pm 0.14 \pm 0.27$	7.3 ± 0.5
$\pi^+\pi^-\pi^0\pi^0$	$10.2 \pm 0.6 \pm 0.7$	---
$\pi^+\pi^+\pi^-\pi^-\pi^0$	$4.31 \pm 0.44 \pm 0.18$	---
$\pi^+\pi^0$	$1.23 \pm 0.06 \pm 0.06$	1.33 ± 0.22
$\pi^+\pi^+\pi^-$	$3.36 \pm 0.10 \pm 0.16$	3.1 ± 0.4
$\pi^+\pi^0\pi^0$	$4.80 \pm 0.27 \pm 0.34$	---
$\pi^+\pi^+\pi^-\pi^0$	$11.7 \pm 0.4 \pm 0.7$	---
$\pi^+\pi^+\pi^+\pi^-\pi^-$	$1.67 \pm 0.18 \pm 0.17$	1.82 ± 0.25
$\eta\pi^+$	$3.56 \pm 0.24 \pm 0.21$	3.0 ± 0.6
$\eta\pi^0$	$0.61 \pm 0.14 \pm 0.05$	---
$\omega\pi^+\pi^-$	$1.66 \pm 0.47 \pm 0.10$	---




Summary of Hadronic Decays

- We understand how to do analyses based on D -tags.
 - We have tuned up the detector simulation to have good data-MC agreement to allow precision measurements.
- The reference modes $D^0 \rightarrow K^- \pi^+$ and $D^+ \rightarrow K^- \pi^+ \pi^+$ are measured at the (1.5 to 2.0)% level in 281 pb^{-1}
 - Systematics dominated – some improvements still possible, but hard to predict how much the systematics can be reduced.
- The extremely clean environment allows us to study basically all D -decays – we should dominate practically all entries in the PDG!


Semileptonic Decays

- Semileptonic decays are easier to describe theoretically
 - The non-perturbative strong physics is parameterized in form factors.
 - For $m_l=0$ we have
 - one form factor for $D \rightarrow (K, \pi) e \nu$
 - three form factors for $D \rightarrow K^* e \nu$
- CLEO-c will measure
 - Inclusive and exclusive branching fractions
 - CKM matrix elements (V_{cd} and V_{cs})
 - Form factors
- The clean environment and excellent detector will allow the first precise studies of Cabibbo suppressed semileptonic D decays
- We only use electrons; muons are too soft to be cleanly identified.



Inclusive Semileptonic D -decays

281 fb⁻¹ (Preliminary)



- This analysis uses only the cleanest tags: $D^0 \rightarrow K^- \pi^+$ and $D^+ \rightarrow K^- \pi^+ \pi^+$
- Correct for e momentum cut
- Obtain the branching fractions

$$Br(D^0 \rightarrow X e \nu_e) = (6.45 \pm 0.17 \pm 0.15) \%$$

$$Br(D^+ \rightarrow X e \nu_e) = (16.19 \pm 0.20 \pm 0.36) \%$$

- Using the measured lifetimes we obtain

$$\frac{\Gamma(D^+ \rightarrow X e \nu_e)}{\Gamma(D^0 \rightarrow X e \nu_e)} = (1.01 \pm 0.03 \pm 0.03)$$

- The sum of exclusive final state

$$\sum_i Br(D^0 \rightarrow X_i e \nu_e) = (6.1 \pm 0.2 \pm 0.2) \%$$



$$\sum_i Br(D^+ \rightarrow X_i e \nu_e) = (15.1 \pm 0.5 \pm 0.5) \%$$

Exclusive Branching Fractions

- The exclusive decays are studied by finding an electron and reconstructing the hadronic system recoiling against a D -tag.
- The signal is extracted by studying the variable $U = E_{\text{miss}} - |P_{\text{miss}}|$
- For the signal events, with one missing neutrino U peaks at zero
- $D \rightarrow K e \nu$ and $D \rightarrow \pi e \nu$ are kinematically separated

CLEO-c 281 pb⁻¹ (Preliminary)






CLEO-III 7 fb⁻¹ (PRL 94:011802, 2005)



Semileptonic D^0 -decays in 56 pb^{-1}






PRL 95, 181802, 2005

First observation




Semileptonic D^+ -decays in 56 pb^{-1}

PRL 95, 181801, 2005



Summary of Exclusive Semileptonic Decays in 56 pb^{-1}





Mode	\mathcal{B} (%)	\mathcal{B} (%) (PDG)
$D^0 \rightarrow \pi^- e^+ \nu_e$	$0.26 \pm 0.03 \pm 0.01$	0.36 ± 0.06
$D^0 \rightarrow K^- e^+ \nu_e$	$3.44 \pm 0.10 \pm 0.10$	3.58 ± 0.18
$D^0 \rightarrow K^{*-}(K^- \pi^0) e^+ \nu_e$	$2.11 \pm 0.23 \pm 0.10$	2.15 ± 0.35
$D^0 \rightarrow K^{*-}(\bar{K}^0 \pi^-) e^+ \nu_e$	$2.19 \pm 0.20 \pm 0.11$	2.15 ± 0.35
$D^0 \rightarrow \rho^- e^+ \nu_e$	$0.19 \pm 0.04 \pm 0.01$	—
$D^+ \rightarrow \pi^0 e^+ \nu_e$	$0.44 \pm 0.06 \pm 0.03$	0.31 ± 0.15
$D^+ \rightarrow \bar{K}^0 e^+ \nu_e$	$8.71 \pm 0.38 \pm 0.37$	6.7 ± 0.9
$D^+ \rightarrow \bar{K}^{*0} e^+ \nu_e$	$5.56 \pm 0.27 \pm 0.23$	5.5 ± 0.7
$D^+ \rightarrow \rho^0 e^+ \nu_e$	$0.21 \pm 0.04 \pm 0.01$	0.25 ± 0.10
$D^+ \rightarrow \omega e^+ \nu_e$	$0.16^{+0.07}_{-0.06} \pm 0.01$	—



- Most modes are improvements over the PDG
 - Including two first observations
 - $D^0 \rightarrow \rho^- e^+ \nu_e$ and $D^+ \rightarrow \omega e^+ \nu_e$
- Most systematics can be reduced with more data
- Updating analysis to 281 pb^{-1}

Exclusive Signals (281 pb⁻¹)

Preliminary



V_{cd} and V_{cs} Determination (281 pb^{-1})

Using LQCD (PRL 94, 011601 (2005))

$$f_+^K(0) = 0.73 \pm 0.03 \pm 0.07 \quad \text{and} \quad f_+^\pi(0) = 0.64 \pm 0.03 \pm 0.06$$

We obtain

$$\blacklozenge D^0 \rightarrow K^- e^+ \nu \quad |V_{cs}| = 0.977 \pm 0.102(\text{LQCD}) \pm 0.009(\alpha) \pm 0.011(\mathcal{B})$$

$$\blacklozenge D^+ \rightarrow K_S e^+ \nu \quad |V_{cs}| = 0.975 \pm 0.102(\text{LQCD}) \pm 0.009(\alpha) \pm 0.015(\mathcal{B})$$

$$\blacklozenge D^0 \rightarrow \pi^- e^+ \nu \quad |V_{cd}| = 0.227 \pm 0.024(\text{LQCD}) \pm 0.006(\alpha) \pm 0.005(\mathcal{B})$$

$$\blacklozenge D^+ \rightarrow \pi^0 e^+ \nu \quad |V_{cd}| = 0.221 \pm 0.023(\text{LQCD}) \pm 0.006(\alpha) \pm 0.009(\mathcal{B})$$

Or using $|V_{cs}| = 0.976 \pm 0.014$ and $|V_{cd}| = |V_{us}| = 0.2252 \pm 0.0008 \pm 0.0021$



$$\blacklozenge D^0 \rightarrow K^- e^+ \nu \quad f_+^K(0) = 0.731 \pm 0.006(\alpha) \pm 0.010(\mathcal{B}) \pm 0.010(V_{cs})$$

$$\blacklozenge D^+ \rightarrow K_S e^+ \nu \quad f_+^K(0) = 0.729 \pm 0.006(\alpha) \pm 0.011(\mathcal{B}) \pm 0.010(V_{cs})$$

$$\blacklozenge D^0 \rightarrow \pi^- e^+ \nu \quad f_+^\pi(0) = 0.645 \pm 0.017(\alpha) \pm 0.014(\mathcal{B}) \pm 0.006(V_{us})$$

$$\blacklozenge D^+ \rightarrow \pi^0 e^+ \nu \quad f_+^\pi(0) = 0.629 \pm 0.017(\alpha) \pm 0.026(\mathcal{B}) \pm 0.006(V_{us})$$

Projections



Form Factors in $D \rightarrow K e \nu$ and $D \rightarrow \pi e \nu$





- The differential decay rate is given by

$$\frac{d\Gamma}{dq^2} = \frac{G_F^2}{24\pi^3} |V_{cq}|^2 p_K^3 |f_+(q^2)|^2$$

- $q^2 = m_{e\nu}^2$
- As we use electrons the form factor corresponding to a scalar W has a negligible contribution to the rate.
- The observed q^2 distribution is fit to extract form factor information.

q^2 Distributions (281 pb⁻¹)

Preliminary



$D \rightarrow (K, \pi) e \nu$ Form Factors

- We fit the observed q^2 distribution to a simple pole form


$$f_+(q^2) = \frac{f_+(0)}{1 - q^2 / M_{\text{pole}}^2}$$

- or a modified pole form (BK)


$$f_+(q^2) = \frac{f_+(0)}{1 - q^2 / M_{D(s)^*}^2} \times \frac{1}{1 - \alpha q^2 / M_{D(s)^*}^2}$$

Form Factor Precision


$D \rightarrow K e \nu$




$D \rightarrow \pi e \nu$



q^2 Dependence in $D \rightarrow \pi e \nu$ (750 pb^{-1})



$D \rightarrow "K^*" e \nu$ Form Factors



- Focus has seen evidence for a S-wave component of the $K\pi$.
- Fit the θ_l and θ_v distributions as a function of q^2 to extract the helicity amplitudes.




$$|A|^2 = \frac{q^2}{8} \left| \begin{array}{l} (1 + \cos \theta_l) \sin \theta_v e^{i\chi} \text{ BW } H_+(q^2) \\ -(1 - \cos \theta_l) \sin \theta_v e^{-i\chi} \text{ BW } H_-(q^2) \\ -2 \sin \theta_l \left(\cos \theta_v \text{ BW } H_0(q^2) + A e^{i\delta} h_0(q^2) \right) \end{array} \right|^2$$

Focus found A 7% of BW peak if they assumed



$$h_0(q^2) = H_0(q^2)$$

$D \rightarrow K^* e \nu$ Form Factor Results

Preliminary




- Non-parametric fit
- Clear evidence for S-wave



Summary of Semileptonic Decays

- CLEO-c measurements are now the best for all semileptonic branching fractions of D mesons
- Form factor studies are now underway
 - Will provide stringent tests of Lattice calculations
- Can measure the form factor normalization – or equivalently the CKM matrix elements.
- In addition to the analyses presented here based on D -tagging, there are neutrino reconstruction style analysis is progress as well.


$$D^+ \rightarrow \mu^+ \nu_\mu \quad \text{and} \quad f_{D^+}$$



$$\Gamma(D^+ \rightarrow l^+ \nu) = \frac{G_F^2}{8\pi} f_{D^+}^2 m_l^2 M_{D^+} \left(1 - \frac{m_l^2}{M_{D^+}^2}\right)^2 |V_{cd}|^2$$


- A precise measurement of f_{D^+} allows precise comparison with theoretical calculations, such as lattice QCD.
- This will help determining f_B , which currently can not be measured in leptonic B decays.

Analysis Technique



- Detect muon and make sure it recoiled against neutrino.
 - Extract signal in M_{miss}^2 which peaks at 0.
- Slightly different use of tags in different analyses

Charged D -tag Reconstruction





Mode	Signal	Background
$K^+ \pi^- \pi^-$	77387 ± 281	1868
$K^+ \pi^- \pi^- \pi^0$	24850 ± 214	12825
$K_S \pi^-$	11162 ± 136	514
$K_S \pi^- \pi^- \pi^+$	18176 ± 255	8976
$K_S \pi^- \pi^0$	20244 ± 170	5223
$K^+ K^- \pi^-$	6535 ± 95	1271
Sum	158354 ± 496	30677

$$M_{BC} = \sqrt{E_{\text{beam}}^2 - |p(D)|^2}$$

Signal Side Selection

- Require one track consistent with coming from the IP for the muon.
 - Muon candidate deposit less than 300 MeV in EM calorimeter
- No additional track from IP
- Require no unmatched showers over 250 MeV
 - Veto background from $D^+ \rightarrow \pi^+ \pi^0$

Highest energy unmatched cluster





Signal Extraction

- For events with μ candidate form

$$MM^2 = (E_{beam} - E_{\mu})^2 - (-\vec{p}_D - \vec{p}_{\mu})^2$$
- Signal will peak at $MM^2 = m_{\nu}^2 = 0$

“Yellow book” MC Study



$D^+ \rightarrow \mu^+ \nu_\mu$ Results

- 50 signal candidate events with the following backgrounds

Background	\mathcal{B} (%)	# of events
$D^+ \rightarrow \pi^+ \pi^0$	0.13 ± 0.02	$1.40 \pm 0.18 \pm 0.22$
$D^+ \rightarrow K^0 \pi^+$	2.77 ± 0.18	$0.33 \pm 0.19 \pm 0.02$
$D^+ \rightarrow \tau^+ \nu$	$2.6 \times \mathcal{B}(D^+ \rightarrow \mu^+ \nu)$	$1.08 \pm 0.15 \pm 0.16$
$D^0 \bar{D}^0, D^+ D^-$	—	$< 0.4, < 0.4, 90\% \text{ C.L.}$
continuum	—	$< 1.2 \text{ } 90\% \text{ C.L.}$
Total		$2.81 \pm 0.30 \pm_{-0.27}^{+0.84}$

- With 158,354 D^+ tags and an efficiency of 67.7% for signal events to satisfy the selection criteria given a D^+ tag we obtain:

$$Br(D^+ \rightarrow \mu^+ \nu) = (4.40 \pm 0.66_{-0.12}^{+0.09}) \times 10^{-4} \quad f_{D^+} = (222.6 \pm 16.7_{-3.4}^{+2.8}) \text{ MeV}$$

(Accepted by PRL)


- We also obtain $Br(D^+ \rightarrow e^+ \nu) < 2.4 \times 10^{-5}$ at 90 C.L.

Systematics $\mathcal{B}(D^+ \rightarrow \mu^+ \nu_\mu)$

	Systematic errors (%)
MC statistics	0.4
Track finding	0.7
PID cut	1.0
MM ² width	1.0
Minimum ionization cut	1.0
Number of tags	0.6
Extra showers cut	0.5
Background	0.6
Total	2.1

- Systematics much smaller than statistical error ($\sim 15\%$)
- At 750 pb^{-1} statistical error 9% on $\mathcal{B}(D^+ \rightarrow \mu^+ \nu_\mu)$
 - or 4.5% on f_{D^+}


Comparing with Theory



D_s Scan

- We performed a scan of the D_s threshold region earlier this fall
 - We took data at point about 20 MeV apart from 4.0 to 4.2 GeV
 - About 5 pb⁻¹ per point in initial scan
 - We analyzed the data within one hour to decide where to run next.
- $D_s D_s$ production at 4.03 GeV was found to be small (0.3 nb)
- Best point is ~4160 MeV with $D_s D_s^*$ cross-section of ~0.9 nb.
 - Additional photon cost ~30% efficiency loss if you have to reconstruct the photon.
- We took about 19 pb⁻¹ of data equivalent to the peak cross-section
 - We have looked at this data...

D_s Branching Fractions



- We have about 36 double tags in the scan data (4140-4200 MeV)
 - About 17% errors on absolute branching fractions
- We will run over the holidays for D_s physics
 - Should have at least 4 times more data for winter conferences

Conclusions

- CLEO-c has accumulated $\sim 281 \text{ pb}^{-1}$ of data at the $\psi(3770)$
- The detector is now well understood and we are producing very precise results based on our current data sample
- We have about 10% D^+ and 17% D^0 tagging efficiency
- CLEO-c has a good start to measure practically all decays that are in the PDG
- Absolute hadronic branching fractions results from 56 pb^{-1} are being updated for 281 pb^{-1} . Statistical precision of $\sim 1\%$.
 - Many other hadronic final states are studied
- Semileptonic and leptonic decays are providing detailed tests of calculations of strong interactions:
 - Form factors for $D \rightarrow (K, \pi) e \nu$
 - Decay constant, f_D , from $D^+ \rightarrow \mu^+ \nu_\mu$
- CLEO-c with 750 pb^{-1} will provide detailed tests of strong interactions
- D_s scan successfully completed, will run at $\sim 4170 \text{ MeV}$

Backup Slides





Comparison to Yellow Book

- Luminosity – lower than we had assumed, discussed earlier.
- Cross-section – we observe ~ 6.4 nb, compare to 10nb assumed.
- Tagging efficiency
 - D^0 Yellow Book had 14% we can use 17%
 - D^+ Yellow Book had 7.5% we can use 10.5%
- Some tags have a higher level of background, but are usable.

Tracking Efficiencies


- For example, we want to measure $B(D \rightarrow K\pi)$ to better than 1%
- We need to measure the tracking efficiency to $\sim 0.3\%$ in order to achieve this goal.
- Luckily, we have data samples that allow crosschecks at this level.
- A very clean sample is the $\psi' \rightarrow J/\psi \pi^+ \pi^-$, with $J/\psi \rightarrow e^+ e^-$ or $\mu^+ \mu^-$.
 - To measure the, *e.g.*, the efficiency for finding a π^+ we reconstruct the J/ψ and the π^- . Then we check if the J/ψ and π^- are consistent with a missing π^+ .
 - This allows us to count the number of events of the type $\psi' \rightarrow J/\psi \pi^+ \pi^-$ without actually finding the π^+ .
 - Now we can simply measure the efficiency by seeing how often we actually find the π^+ in the event.

Pion Tracking Efficiency



Rare D -decays

- CLEO-c has searched for the rare decays $D \rightarrow (K, \pi)e^+e^-$.
- No significant signal observed, limits a factor of 10 better than PDG.



- Searches also underway for $D \rightarrow \phi\gamma$ and $D \rightarrow K^*\gamma$
- These rare decays are likely to be long distance dominated, but one should still look for them.